

Yang-Baxter Algebra for the n -Harmonic Oscillator Realisation of $sp(2n, \mathbb{R})$

A.J. Macfarlane and F. Wagner

D.A.M.T.P

Silver Street, Cambridge CB3 9EW, UK

February 9, 2008

Abstract

Using a rational R -matrix associated with the 4×4 defining matrix representation of $c_2 \cong sp(4)$, the Lie algebra of $Sp(4)$, a one-site operator solution of the associated Yang-Baxter algebra acting in the Fock space of two harmonic oscillators is derived. This is used to define N -site integrable systems, which are soluble by a version of the algebraic Bethe ansatz method without nesting. All essential aspects of the work generalise directly from c_2 to c_n .

1 Introduction

One can see from references such as [1] that the algebraic Bethe ansatz [2] continues to be a subject of interesting development and improvement. This paper is devoted to applying the method to integrable models with underlying Lie algebras other than $a_1 \cong su(2)$ in a way that produces *unnested* Bethe equations.

Let $\mathcal{V}$ and $\mathcal{H}$ denote auxiliary and quantum spaces respectively. Define $R(u)$ with spectral parameter u in $\mathcal{V} \otimes \mathcal{V}$, and $T(u)$ in $\mathcal{V} \otimes \mathcal{H}$. Then the Yang-Baxter equation for $R(u)$ on $\mathcal{V} \otimes \mathcal{V} \otimes \mathcal{V}$,

$$R_{12}(u-v)R_{13}(u)R_{13}(v) = R_{23}(v)R_{13}(u)R_{12}(u-v) \quad , \quad (1)$$

and the Yang-Baxter operator algebra relation on $\mathcal{V} \otimes \mathcal{V} \otimes \mathcal{H}$,

$$R_{12}(u-v)T_1(u)T_2(v) = T_2(v)T_1(u)R_{12}(u-v) \quad , \quad (2)$$

are objects of central importance to the study of integrable systems. In contexts in which there is an underlying algebra of type a_n , especially a_1 , things are quite well understood. See *e.g.* [3, 4, 5]. However, even for a_n for $n > 1$, solution of related models by the Bethe ansatz method is iterative, giving eigenstates specified by means of nested Bethe equations [6]. The same applies also to models with $so(n)$ or $c_n \cong sp(2n)$ as underlying Lie algebra [7, 8]. A paper that describes the nested Bethe ansatz method clearly is [9]. Papers which state the case for seeking, and which implement, alternatives to that method with its evident complications are [10] (on Gaudin models) and [11].

Here we deal with theories governed by $c_n \cong sp(2n)$. We first obtain a solution of (1), and then a new ‘one-site’ solution $L(u) = T(u)$ of (2) in $\mathcal{V} \otimes \mathcal{H}$ when $\mathcal{H}$ is the representation space of the metaplectic representation $\mathcal{M}_n$ of c_n defined in terms of $2n$ pairs of harmonic oscillator variables. This is the main result of the paper. Given this

$L(u)$ however, we can proceed directly down the path laid down by Sklyanin [4] for the construction of families of N -site integrable models.

The integrable models just indicated are of interest because they are soluble by means of a one stage application of the Bethe ansatz method, giving rise to Bethe equations without any nesting, equations that are no more complicated than those familiar within a_1 studies [5].

In this paper, we present some essential results for c_2 and indicate their extension to c_n in section 2, describe $\mathcal{M}_n$ in section 3, and derive our formulas for $R(u)$ and $L(u)$ in section 4. Section five contains a sketch of the solution of our c_2 integrable models by our Bethe ansatz method. Fuller discussion of this last topic and various related matters will be provided in papers in preparation [12].

2 The Lie algebra c_n

We present the basis of c_2 in Cartan-Weyl form with simple roots $\mathbf{r}_1 = (1, -1)$, $\mathbf{r}_2 = (0, 2)$, so that the other positive roots are $\mathbf{r}_3 = \mathbf{r}_1 + \mathbf{r}_2 = (1, 1)$ and $\mathbf{r}_4 = \mathbf{r}_1 + \mathbf{r}_3 = (2, 0)$. Let $\mathbf{H} = (H_1, H_2)$ denote the generators of the Cartan subalgebra of c_2 . Let $E_{\pm\alpha}$, $\alpha \in \{1, \dots, 4\}$, denote the raising and lowering generators. Then the Lie algebra c_2 is specified by

$$\begin{aligned} [H_i, E_{\pm\alpha}] &= \pm(\mathbf{r}_\alpha)_i E_{\pm\alpha} \quad , \quad [E_\alpha, E_{-\alpha}] = \mathbf{r}_\alpha \cdot \mathbf{H} \quad , \\ [E_1, E_2] &= \sqrt{2}E_3 \quad , \quad [E_1, E_3] = \sqrt{2}E_4 \quad , \quad \text{etc.} \end{aligned} \quad (3)$$

We also use a Cartesian basis X_i , $i \in \{1, \dots, 10\}$, for c_2

$$H_i = X_i \quad , \quad H_2 = X_2 \quad \sqrt{2}E_{\pm\alpha} = X_{2\alpha+1} + iX_{2\alpha+2} \quad . \quad (4)$$

Then we may define the structure constants of c_2 via

$$[X_i, X_j] = ic_{ijk}X_k \quad , \quad (5)$$

and the quadratic Casimir is

$$\mathcal{C}^{(2)} = X_i X_i = H_1^2 + H_2^2 + \sum_{\alpha} \{E_{\alpha}, E_{-\alpha}\} \quad . \quad (6)$$

We also require the matrices of the four-by-four defining representation $\mathcal{D}$ of c_2 , defined according to

$$X_i \mapsto x_i \quad ; \quad H_1 \mapsto h_1 \quad , \quad H_2 \mapsto h_2 \quad , \quad E_{\pm\alpha} \mapsto e_{\pm\alpha} \quad . \quad (7)$$

We can display these all at once by use of a 4×4 matrix $C = x_i \otimes X_i \equiv x_i X_i$ that is very useful for work on the c_2 Yang-Baxter algebra. We have

$$\begin{aligned} C &= H_1 h_1 + H_2 h_2 + \sum_{\alpha=1}^4 (E_{\alpha} e_{-\alpha} + E_{-\alpha} e_{\alpha}) \\ &= \begin{pmatrix} H_1 & E_{-1} & E_{-3} & \sqrt{2}E_{-4} \\ E_1 & H_2 & \sqrt{2}E_{-2} & E_{-3} \\ E_3 & \sqrt{2}E_2 & -H_2 & -E_{-1} \\ \sqrt{2}E_4 & E_3 & -E_1 & -H_1 \end{pmatrix} \quad . \end{aligned} \quad (8)$$

One may read the explicit forms for $h_1, h_2, e_{\pm\alpha}$ off (8), verify that they do represent (3) correctly, and note the hermiticity properties $h_1^\dagger = h_1, h_2^\dagger = h_1, e_{\pm\alpha}^\dagger = e_{\mp\alpha}$ appropriate to generators of the compact representation $\mathcal{D}$ of c_2 . It can further be seen that the matrices $x_i, i \in \{1, 2, \dots, 10\}$, possess the properties

$$x_i^\dagger = x_i, \quad \text{Tr}(x_i) = 0, \quad \text{Tr}(x_i x_j) = 2\delta_{ij}, \quad x_i^T = -J x_i J^T \quad . \quad (9)$$

Here J , the standard c_2 symplectic form, in our representation, has, as its only non-zero elements: $J_{14} = J_{23} = -J_{32} = -J_{41} = 1$. In view of (9), it follows that the matrices x_i satisfy the completeness relation

$$x_{iac} x_{ibd} = \delta_{ad} \delta_{bc} - J_{ab} J_{cd} \quad \text{or} \quad x_i \otimes x_i = P - K \quad , \quad (10)$$

where $K_{ab,cd} = J_{ab} J_{cd}$, is proportional to the projector onto the symplectic trace, and P is the permutation map on $\mathbb{C}^4 \otimes \mathbb{C}^4$.

In the defining representation $X_i \mapsto x_i$ the Casimir $\mathcal{C}^{(2)}$ has eigenvalue 5. In the adjoint representation $(X_i)_{jk} = -i c_{ijk}$, it has eigenvalue 12, so that also

$$c_{ijk} c_{ijl} = 12 \delta_{kl} \quad . \quad (11)$$

Our work on the Yang-Baxter algebra requires us to extend the span of the x_i to the space of all traceless hermitian 4×4 matrices. Thus we define a specific set of five matrices $y_a, a \in \{1, 2, \dots, 5\}$, with the properties

$$y_a^\dagger = y_a, \quad \text{Tr} y_a = 0, \quad \text{Tr} y_a y_b = 2\delta_{ab}, \quad y_a^T = J y_a J^T \quad , \quad (12)$$

so that the $J x_i$ define a set of 10 symmetric matrices, whereas $J y_a$ and J itself define a set of 6 antisymmetric ones. Further we can write

$$x_i x_j + x_j x_i = \delta_{ij} + 2 d_{ija} y_a \quad , \quad (13)$$

which defines $d_{ija} = d_{jia}$ such that $d_{iia} = 0$. Use of the completeness relation (9) and trace properties, allows proof of the results

$$d_{ija} d_{ijb} = 3\delta_{ab} \quad , \quad y_a = \frac{1}{3} d_{ija} x_i x_j \quad . \quad (14)$$

Our definition (14) of the y_a reflects the fact that the set $\lambda_A = \{x_1, \dots, x_{10}, y_1, \dots, y_5\}$ can serve as a set of 15 a_3 Gell-Mann λ -matrices which satisfy

$$\lambda_A^\dagger = \lambda_A, \quad \text{Tr} \lambda_A = 0, \quad \text{Tr} \lambda_A \lambda_B = 2\delta_{AB}, \quad \lambda_A \otimes \lambda_A = 2P - \frac{1}{2} I_4 \quad , \quad (15)$$

where I_4 is the 4×4 unit matrix. If we write $X = x_i \otimes x_i, Y = y_a \otimes y_a$, then the last part of (15) and (10) read as

$$X + Y = 2P - \frac{1}{2} I_4, \quad X = P - K \quad , \quad (16)$$

which can be solved for P and K in terms of I_4, X, Y .

3 The Metaplectic Representation, $\mathcal{M}_n$ of c_n .

Just as the metaplectic representation of $c_1 \cong a_1$ is discussed using the Fock space of one harmonic oscillator [13, 14] so also is that of c_n discussed using n independent oscillators.

We present results explicitly for c_2 in a form that generalises completely and naturally for c_n for all n .

Since we define the row vector v^T in terms of two sets of standard harmonic oscillator variables by

$$v^T = (a_1^\dagger \ a_2^\dagger \ a_2 \ a_1) \quad , \quad (17)$$

then, using also the definition of J given above, it can be shown explicitly that the operators

$$X_i = \frac{1}{2} v^T x_i J v \quad (18)$$

obey the same commutation relations (5) as the matrices x_i of (7). Then (4) leads us to the explicit results

$$\begin{aligned} H_1 &= \frac{1}{2} \{a_1, a_1^\dagger\}, \quad H_2 = \frac{1}{2} \{a_2, a_2^\dagger\} \quad , \quad E_{+1} = a_1^\dagger a_2, \quad E_{-1} = a_1 a_2^\dagger, \\ E_{+2} &= -a_2^{\dagger 2} / \sqrt{2}, \quad E_{+3} = -a_1^\dagger a_2^\dagger, \quad E_{+4} = -a_1^{\dagger 2} / \sqrt{2}, \\ E_{-2} &= a_2^2 / \sqrt{2}, \quad E_{-3} = a_1 a_2, \quad E_{-4} = a_1^2 / \sqrt{2}. \end{aligned} \quad (19)$$

One can also check directly that (19) correctly provides a representation of (3). From (19), we find the hermiticity conditions $H_1^\dagger = H_1$, $H_2^\dagger = H_2$, $E_{+1}^\dagger = E_{-1}$, $E_\alpha^\dagger = -E_{-\alpha}$, $\alpha \in \{2, 3, 4\}$. These reflect the fact that (19) generates the infinite dimensional unitary metaplectic representation $\mathcal{M}_2$ of the real non-compact groups $Sp(2, \mathbb{R})$. Also the maximal compact subalgebra of (19) is that of an $SU(2) \times U(1)$ group, with angular momentum type generators given by

$$J_z = \frac{1}{2} (H_1 - H_2), \quad J_+ = a_1^\dagger a_2 = E_{+1} = J_-^\dagger, \quad J_- = a_2^\dagger a_1 = E_{-1} \quad (20)$$

and $U(1)$ generator $\lambda = \frac{1}{2} (H_1 + H_2)$, $[\lambda, \vec{J}] = \mathbf{0}$. In addition, we note the two commuting a_1 subalgebras generated by $H_1, E_{\pm 4}$ and $H_2, E_{\pm 2}$. These correspond to the two c_1 metaplectic subrepresentations of our c_2 metaplectic representation (19).

We also need to verify that the ‘non-compact’ raising and lowering generators of c_2 can be arranged to define vector operators $\vec{\xi}$ and $\vec{\chi}$, using the standard Racah definition of tensor operators. Explicitly $\vec{\xi}$ and $\vec{\chi}$ have spherical components

$$(\xi_{-1}, \xi_0, \xi_{+1}) = (E_2, E_3, E_4) \quad , \quad (\chi_{-1}, \chi_0, \chi_{+1}) = (E_{-4}, -E_{-3}, E_{-2}) \quad . \quad (21)$$

All the required properties can be checked explicitly. It is significant for the Bethe ansatz solution of the integrable models we construct, that the components of the vectors ξ commute with each other, as do those of χ .

We are now in a position to derive the key algebraic result for $\mathcal{M}_2$, a result which allows us to obtain an infinite dimensional operator solution of (1) when $\mathcal{M}_2$ is taken as the quantum space $\mathcal{H}$, (and similarly for c_n). To this end we use (18) and (10) to write the matrix $C = x_i X_i$ of (4) in the form

$$C = -\frac{1}{2} I_4 + (Jv)v^T \quad , \quad (22)$$

whence it follows that C obeys the quadratic relation

$$C^2 = -3C - \frac{5}{4} I_4 \quad . \quad (23)$$

We stress that this non-trivial result holds for the particular (metaplectic) representation (19) of c_2 . It does not hold in an arbitrary representation of c_2 .

Since $\text{Tr } C = 0$, (23) implies that for $\mathcal{M}_2$ we have $\mathcal{C}^{(2)} = \frac{1}{2}\text{Tr } C^2 = -\frac{5}{2}$, a result that can be checked directly to be correct by inserting (19) into (6), and expressing everything in terms of $N_1 = a_1^\dagger a_1$, $N_2 = a_2^\dagger a_2$.

The entire discussion generalises readily from c_2 to c_n . The generalisation of (18), and of the completeness relations (12) are obvious, and imply that the only change needed in (22) replaces the unit matrix I_4 of $\mathbb{C}^4$ by I_{2n} . Then we have

$$C^2 = -(n+1)C - \left(\frac{2n+1}{4}\right) I_{2n} \quad , \quad (24)$$

so that

$$\mathcal{C}^{(2)} = \frac{1}{2}\text{Tr } C^2 = -\frac{1}{4}\text{dim}(c_n). \quad (25)$$

4 The Yang-Baxter Equation and Algebra of c_n

Let $\mathcal{V} = \mathbb{C}^4$. Then we seek a solution of the Yang-Baxter equation (1) of c_2 of the form $R(u) = uI + \eta P + C(u)K$, or

$$R_{ab,cd}(u) = u\delta_{ac}\delta_{bd} + \eta\delta_{ad}\delta_{cb} + C(u)J_{ab}J_{cd} \quad . \quad (26)$$

This is a natural generalisation of the work for a_{2n} where only I and P are invariant under the action of $sl(2n, \mathbb{C})$, to the present case where K also is invariant under the action of $sp(2n, \mathbb{C})$.

Equation (26) yields a solution of (1) provided that $C(u)$ is given by

$$C(u) = \frac{u\eta}{\lambda - u}, \quad \lambda = -3\eta \quad , \quad (27)$$

and the same holds for c_n when $\lambda = -(1+n)\eta$. These results are the symplectic analogues of results found for the orthogonal groups, *e.g.* in [15], and agree with results presented in different form, *e.g.* in [3]. We wish to use the R -matrix (26), (27) to reach solutions of the Yang-Baxter algebra on $\mathcal{V} \otimes \mathcal{V} \otimes \mathcal{H}$ when $\mathcal{H}$ is the module corresponding to the operator representation given by (19).

First as a guide we review the route followed successfully and in generality for the same purpose in the case of the a_n algebras. There one has

$$R(u) = uI + \eta P = uI + \eta \left(\frac{1}{2} \sum_A \lambda_A \otimes \lambda_A + \frac{1}{n} I \right) \quad , \quad (28)$$

using the completeness relation of the Gell-Mann λ -matrices of a_n . Changing the representation $\frac{1}{2}\lambda_A$ to X_A leads to

$$L(u) = (uI + \frac{1}{n}\eta) + \eta \sum_A \lambda_A \otimes X_A \quad . \quad (29)$$

which can be seen to give a ‘one-site’ solution $T(u) = L(u)$ of (2).

The same approach does not work straightforwardly for c_2 (or c_n) because (26) involves not only $X = x_i \otimes x_i$ but also $Y = y_a \otimes y_a$, where (*cf.* section 2) the y_a do not belong to

the Lie algebra c_2 . However, as we have shown above, they can be expressed in terms of the x_i and so we enforce the change of representation from x_i to X_i onto Y

$$\begin{aligned} y_a \otimes y_a &= y_a \otimes \frac{1}{3} d_{ija} x_i x_j \mapsto y_a \otimes \frac{1}{3} d_{ija} X_i X_j \\ &= \frac{1}{6} (-\delta_{ij} + \{x_i, x_j\}) \otimes X_i X_j = \frac{1}{3} (C^2 + 3C - \frac{1}{2} \mathcal{C}^{(2)}) \quad . \end{aligned} \quad (30)$$

where, as before, $C = x_i \otimes X_i \equiv x_i X_i$, $\mathcal{C}^{(2)} = X_i X_i$. The results (14), (13) and (5) have all been used here. We do not know how to handle the complication that follows the general use of (30), but we do know how to proceed in special cases, such as that of section 3, in which $\mathcal{C}^{(2)} = -\frac{5}{2}$ and (23) holds, for then the RHS of (30) vanishes. Accordingly we seek a solution $T(u)$ or first $L(u)$ of (2) in the form

$$L(u) = \alpha(u)I + \beta(u)x_i X_i \quad , \quad (31)$$

This ansatz succeeds for

$$\frac{\alpha(u)}{\beta(u)} = \frac{u - \delta}{\eta}, \quad \delta \in \mathbb{R} \quad , \quad (32)$$

and we may put $\beta(u) = \eta$ and $\alpha(u) = u - \delta$. We emphasise the non-trivial nature of the result obtained. Since $R(u)$ given by (26) necessarily involves $Y = y_a \otimes y_a$ the ansatz (31) is not obviously valid a priori in our work. It seems unlikely to be valid for general representations in the quantum space. But it works for metaplectic representation of c_2 , and likewise for other c_n .

5 Bethe Ansatz for c_2 Integrable Models

Sklyanin's general procedure [4] for constructing integrable spin models, *e.g.* if operators $X_i = S_i$ acting in $\mathcal{H} = \mathcal{V}_s$ where $\mathcal{V}_s$ is the vector space in which the spin- s representation of $SU(2)$ acts, is available here also once we know a basic ('one-site') matrix $L(u) \in \mathcal{V} \otimes \mathcal{H}$ which satisfies (2). We write

$$T(u) = \mathcal{K} \prod_{r=1}^N L(u - \delta_r) \quad , \quad (33)$$

where $\mathcal{K}$ is a constant matrix such that $[\mathcal{K} \otimes \mathcal{K}, R(u)] = 0$. Since $T(u)$ is an n -site solution of (2), it follows in a well-known way that $[t(u), t(v)] = 0$ where $t(u) = \text{Tr } T(u)$, and expansion of $t(u)$ in powers of u yields constants of the motion.

In this section we sketch the solution for the eigenstates of $t(u) = \sum_{i=1}^4 T_{ii}$ of the system described by (33) when $L(u)$ is the one-site solution of (2) given in the last section by means of the algebraic Bethe ansatz method [2, 4]. Although detailed attention to $\mathcal{K}$ and to the δ_r is important for the discussion of the completeness of the set of these eigenstates, we defer this to a later publication, here setting $\mathcal{K} = 1$ and $\delta_r = 0$. We wish rather to explain how the Bethe ansatz method works in our models, emphasising the simplifications that stem from special features of the representation (19) of X_i , and exploiting the transformation properties of the T_{ij} under the compact $su(2)$ subgroup of c_2 . The latter follow for the T_{ij} when $\mathbf{J} = \sum_{r=1}^N \mathbf{J}_r$ is used, J_r for $r = 1, \dots, N$ being defined at the r -th site in terms of the oscillator variables of that site as in (19). Perhaps it should be pointed out that $\mathbf{J}$ as just defined is not equal to the $(N-1)$ -fold coproduct of the 'one-site' operator ($J_+ = T_{21}^{(1)}$, $J_z = \frac{1}{2}(T_{11}^{(1)} - T_{22}^{(1)})$, $J_- = T_{12}^{(1)}$), in the sense of

the Yang-Baxter algebra (2). Nevertheless its commutation relations with the $\Delta^{(N)}T_{ij}^{(1)}$, involving the standard comultiplication, can be shown explicitly to give these operators the correct $su(2)$ tensor operator properties.

Our sketch deals only with the tower of Bethe states based in the Fock vacuum of both the oscillators at each of the N sites. Similar work for other towers whose ground states involve oscillator states of occupation number one requires only a modest extension of the analysis described here. See [12]. We note first, from (19), that all the L_{ij} with $i < j$, as well as L_{21} and L_{43} annihilate the Fock vacuum at each site, and it is easy to promote the same result to the same set of N -site T_{ij} of (33). Also, at any site we have $(L_{42} - L_{31})|0\rangle = 0$, and this result can be similarly promoted to

$$(T_{42} - T_{31})(u)|0\rangle = 0 \quad . \quad (34)$$

We aim first to construct the Bethe states

$$T_{41}(v)|0\rangle \quad , \quad T_{42}(v_1)T_{31}(v_2)|0\rangle \quad , \quad \dots \quad , \quad (35)$$

which have $j = m = 1, 2, \dots$ quantum numbers w.r.t $\mathbf{J}$. Then the other states with the same j and lower m follow by application of the lowering operator J_- . Since $[\mathbf{J}, t(u)] = 0$ all the states of each multiplet of Bethe states have the same eigenvalue of $t(u)$, and allowed v -values given by the same set of Bethe equations. One naturally expects (and finds) that there are, alongside the $j = 2$ multiplet, $j = 1$ and $j = 0$ multiplets bilinear in the creation operators T_{41}, T_{31}, T_{42} and T_{32} . There is not scope in this paper to describe in full the subtleties of the analysis (in which (34) allows significant simplifications), so next we present results.

Let $w = u - v$, let $\tau_+(u)$ denote the eigenvalue of T_{11} and of T_{22} for the vacuum state $|0\rangle$, and let $\tau_-(u)$ do the same for T_{33} and T_{44} . It is easy to calculate $\tau_{\pm}(u)$ using (33) and (19). Then for $j = 1$, we find

$$t(u)T_{41}(v)|0\rangle = \tau(u)T_{41}(v)|0\rangle \quad , \quad (36)$$

where the eigenvalue $\tau(u)$ is given by

$$\tau(u) = (1 + \frac{\eta}{w})\tau_+(u) + (1 - \frac{\eta}{w})\tau_-(u) \quad , \quad (37)$$

and v is determined by the Bethe equations $\tau_+(v) = \tau_-(v)$.

For $j = m = 2$, the eigenvalue $\tau(u)$ of $t(u)$ for $T_{41}(v_1)T_{41}(v_2)|0\rangle$, and the Bethe equations for v_1 and v_2 are given by

$$\begin{aligned} \tau(u) &= 2(1 + \frac{\eta}{w_1} + \frac{\eta}{w_2} + 2\frac{\eta^2}{w_1w_2})\tau_+(u) \\ &+ 2(1 - \frac{\eta}{w_1} - \frac{\eta}{w_2} + 2\frac{\eta^2}{w_1w_2})\tau_-(u) \quad , \end{aligned} \quad (38)$$

where we write $w_k = u - v_k$, $k = 1, 2$ and $v = v_1 - v_2$, and

$$\tau_+(v_1)/\tau_-(v_1) = \tau_-(v_2)/\tau_+(v_2) = (v + 2\eta)/(v - 2\eta) \quad . \quad (39)$$

The generalisation to higher integral j -values is evident. Turning to $m = 1$ states bilinear in the creation T_{ij} , we find two, one that follows by application of J_- to the state just discussed. Writing

$$\begin{aligned} \phi_1 &= T_{42}(v_1)T_{41}(v_2)|0\rangle \quad , \\ \phi_2 &= T_{31}(v_1)T_{41}(v_2)|0\rangle \quad , \\ \phi_3 &= T_{41}(v_1)T_{42}(v_2)|0\rangle \quad , \\ \phi_4 &= T_{41}(v_1)T_{31}(v_2)|0\rangle \quad , \end{aligned} \quad (40)$$

we find the $|21\rangle$ state is given by $(\phi_1 + \phi_2 + \phi_3 + \phi_4)$, and the unique $|11\rangle$ Bethe state in the context is given by $\{(\phi_1 + \phi_2 - \phi_3 - \phi_4) + \frac{2\eta}{v}(\phi_1 - \phi_2)\}$ with $t(u)$ eigenvalue given by $2(1 + \frac{\eta}{w_1} + \frac{\eta}{w_2})\tau_+(u) + 2(1 - \frac{\eta}{w_1} - \frac{\eta}{w_2})\tau_-(u)$, and v_1 and v_2 determined by the Bethe equations $\tau_+(v_1) = \tau_-(v_1)$, $\tau_+(v_2) = \tau_-(v_2)$. The pattern of the Bethe states should be clear now.

The states constructed have various nice properties. Their Bethe equations ensure that the poles of their $t(u)$ eigenvalues have residues zero. They are also invariant under the exchange of v_1 and v_2 , although proof of this is not obvious, requiring detailed use of the commutation relations of the T_{ij} that we are using.

A brief sketch of how these results are derived is called for, because there is a level of complication not seen in Bethe ansatz studies in models with $a_1 = su(2)$ invariance. This is no doubt the price to be paid for obtaining results without recourse to a nesting process. Complications not present in a_1 studies come into play here whenever the third term of (26) enters the commutation relations amongst the T_{ij} non-trivially. A sufficient illustration of what has to be done whenever this happens is provided by showing how to compute the effect of the term $T_{11}(u)$ of $t(u)$ on $T_{41}(v)|0\rangle$. We here shall abbreviate any $T_{ij}(u)$ or $T_{ij}(v)$ by T_{ij} or T'_{ij} . Our strategy is push to the right end of any term only factors that annihilate $|0\rangle$, or else are of the type T_{kk} , with no sum on k , which multiply $|0\rangle$ by $\tau_{\pm}$. Direct calculation of $wT_{11}T'_{41}$ from (2) fails to achieve this, leaving a term in $T_{21}T'_{31}$ that has to be eliminated by calculation from (2) of $wT_{21}T'_{31}$. This two stage procedure yields finally the result

$$T_{11}T'_{41} = (1 + \frac{2\eta}{w})T'_{41}T_{11} - \frac{2\eta}{w}T_{41}T'_{11} + \frac{\eta}{w}(T'_{31}T_{21} - T_{31}T'_{21}) \quad . \quad (41)$$

Another example yields

$$\begin{aligned} T_{11}T'_{42} &= (1 + \frac{\eta}{w + \eta})T'_{42}T_{11} + (\frac{2\eta}{w} - \frac{\eta}{w + \eta})T'_{41}T_{12} - \frac{2\eta}{w}T_{41}T'_{12} \\ &\quad - \frac{\eta}{w}T_{31}T'_{22} + (\frac{\eta}{w} - \frac{\eta}{w + \eta})T'_{31}T_{22} + \frac{\eta}{w + \eta}T'_{32}T_{21} \quad . \end{aligned} \quad (42)$$

One must push each of the four pieces of $t(u)$ past each of the creator T_{ij} , needing a total of about 32 results in all, to deduce results such as those just quoted. This entails eight results like each of (41) and (42); the remainder do not involve third term of (26) and are written down directly.

6 Discussion

We have described, for c_2 , formalism and the construction of a family of integrable models that generalise in every respect to all c_n . This applies to the construction (18) and the proof that the X_i obey the same commutation relations as the x_i , to the deduction of relations (31) and (32). A similar construction is available for the d_n series, where we use fermionic instead of bosonic oscillator variables. Note that in the c_n case, there is a maximal compact $u(n)$ subalgebra, and sets of creation operators T_{ij} which transform according to its dimension n defining representation $(1, \dots, 0)$, leading to a $u(n)$ multiplet structure of Bethe eigenstates [12], with a pattern similar to that indicated in section five. We note also that our method of getting Bethe eigenstates without recourse to nesting applies equally well to the treatment [12] of c_n Gaudin models [16].

Acknowledgement. This work was partly supported by PPARC. F.W. is grateful to Trinity College, Cambridge for an IGS, and to EPSRC for a research grant. We thank Hendryk Pfeiffer for useful discussions.

References

- [1] A.G. Izergin, N. Kitanine, J.M. Maillet and V. Terras. *Spontaneous magnetization of the XXZ Heisenberg spin chain*. Preprint Lyon, France, LPENSL-TH-13-98.
- [2] L.D. Fadde'ev. *Recent advances in field theory and statistical mechanics*. Les Houches Lectures, eds. J.B. Zuber and R. Stora, Elsevier, 1984, pp561-608.
- [3] M. Jimbo. *Introduction to the Yang-Baxter equation*. Internat. J. Phys. **4** 3759-3777 (1989).
- [4] E.K. Sklyanin. *Separation of variables: new trends*. Prog. Theor. Phys. Supp.. **118** 35-60 (1995).
- [5] C. Gomez, M. Ruiz-Altaba and G. Sierra. *Quantum Groups in two dimensional physics*. Cambridge University Press, 1996.
- [6] N. Yu. Reshitikhin. *Calculation of the norms of Bethe Vectors for $SU(3)$ symmetry*. Zap. Nauchnyk. Sem. LOMI **150** 196-213 (1986) or J. Sov. Math. **46** 1694-1706 (1989).
- [7] N. Yu. Reshitikhin. *Algebraic Bethe ansatz for $SO(N)$ invariant transfer matrices*. Zap. Nauchnyk. Sem. LOMI **169** 128-140 (1988) or J. Sov. Math. **54** 940-951 (1991).
- [8] N. Yu. Reshitikhin. *Integrable models of quantum magnets*. TMF **63** 347-366 (1985) or Theor. Math. Phys. **63** 555-569 (1985).
- [9] O. Babelon, H.J. de Vega and C.M. Viallet. *Exact solution of the $Z_{n+1} \otimes Z_{n+1}$ symmetric generalisation of the XXZ model*. Nucl. Phys. **B200** 266-280 (1982).
- [10] B. Feigin, E. Frenkel and N. Yu. Reshitikhin. *Gaudin model, Bethe ansatz and critical level*. Commun. Math. Phys. **166** 27-62 (1994).
- [11] H.M. Babujian and R. Flume. *Off-shell Bethe ansatz equations and N -point conformal correlators in the $SU(2)$ WZNW theory*. Mod. Phys. Lett. **A9** 2029-2040 (1994).
- [12] A.J. Macfarlane, Hendryk Pfeiffer and F. Wagner. DAMTP Cambridge preprints, in preparation.
- [13] C. Itzykson. *Remarks on the bosonic commutation rules*. Commun. Math. Phys. **4** 92-122 (1967).
- [14] S. Sternberg and J.A. Wolf. *Hermitian Lie algebras and metaplectic representations*. Trans. Am. Math. Soc. **238** 1-43 (1976).
- [15] P.P. Kulish and E.K. Sklyanin. *Solutions of the Yang-Baxter equation*. Zap. Nauchnyk. Sem. LOMI **95** 129-160 (1980) or J. Sov. Math. **19** 1596-1620 (1982).
- [16] M. Gaudin. *Diagonalisation d'une classe d'Hamiltonians de spin*. J. Physique **37** 1087-1098 (1976); *La fonction d'onde de Bethe*. Masson 1983.